

Research Article

Adipose Derived Mesenchymal Stem Cells Efficiently Rescue Carbon Tetrachloride-Induced Acute Liver Failure in Mouse

Lisi Deng,^{1,2} Guangze Liu,¹ Xin Wu,^{1,3} Yaping Wang,⁴ Minghua Tong,¹ Bing Liu,¹ Kuidong Wang,¹ Yuting Peng,^{1,3} and Xiangping Kong¹

¹ Center of Infectious Diseases, 458th Hospital of PLA, No. 801 Dongfengdong Road, Guangzhou, Guangdong 510600, China

² Southern Medical University, Guangzhou, Guangdong 510515, China

³ Institute of Life Science and Bio-Pharmaceutics, Shenyang Pharmaceutical University, Shenyang 110016, China

⁴ Guangzhou Eight People's Hospital, Guangdong 510060, China

Correspondence should be addressed to Xiangping Kong; xiangping_kong@hotmail.com

Received 22 January 2014; Accepted 9 May 2014; Published 4 June 2014

Academic Editor: Vito Pistoia

Copyright © 2014 Lisi Deng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background and Aim. Adipose derived mesenchymal stem cells (ADMSCs) may be an attractive source for acute and chronic liver injury because they are abundant and easy to obtain. We aim to investigate the efficacy of ADMSCs transplantation in the acute liver failure (ALF) caused by carbon tetrachloride (CCl_4) in mice. **Methods.** ADMSCs were isolated from inguinal fat pads of enhanced green fluorescent protein (EGFP) transgenic mice and their surface markers and differentiation potential were analyzed. ALF models were established by infusion of CCl_4 and divided into two groups: control group; EGFP-ADMSCs transplantation group. The restoration of biological functions of the livers receiving transplantation was assessed via a variety of approaches such as survival rates, live function parameters, histological localization of EGFP-ADMSCs, and Immunofluorescence analysis. **Results.** ADMSCs were positive for CD105, CD44 but negative for CD45, CD34 and had adipogenic, osteogenic differentiation potential. The survival rate of transplantation group significantly increased compared to PBS group. Furthermore, the transplanted cells were well integrated into injured livers and produced albumin, cytokeratin-18. **Conclusion.** Direct transplantation of ADMSCs is an effective treatment for ALF. The transplanted ADMSCs exhibit the potential to differentiate into hepatocyte-like cells in the injured livers.

1. Introduction

Acute liver failure (ALF), with a high mortality rate, is a serious clinical condition. Up to now, orthotopic liver transplantation is the only resolutive treatment for ALF and end-stage liver disease [1]. However, for the reason of limited number of donors and organ rejection, alternative approaches are needed. Currently, the development of cell therapy for the treatment of end-stage hepatic diseases is under active investigation [2, 3]. And the cell therapy can be defined as “the use of living cells to restore, maintain, or enhance tissue and organ function” [4]. While the preeminent candidate stem cells for therapy of an injured liver are mesenchymal stem cells (MSCs), MSCs are adherent, fibroblast-like, pluripotent, and nonhematopoietic progenitor cells, which reside in various tissues such as the bone marrow [5], adipose tissue [6], umbilical cords [7], and placenta [8]. Adipose derived

mesenchymal stem cells (ADMSCs) are an attractive source for regenerative medicine because they are abundant and easy to obtain from liposuction aspirates or excised fat [9]. They also display multilineage differentiation potential and can differentiate into hepatocyte-like cells *in vitro* [10, 11]. Moreover, ADMSCs are well tolerated and exert *in vitro* and *in vivo* immunoregulatory properties [12]. It has been reported that ADMSCs demonstrate therapeutic efficacy in liver injuries and secrete cytokines and growth factors associated with hepatic regeneration [13–17]. However, the use of ADMSCs to treat acute liver failure in animal models is largely unknown. Additionally, because of difficulty in tracking transplanted ADMSCs-derived hepatocytes in patients, the roles of ADMSCs in liver regeneration have not been fully elucidated. In the current study, we investigated whether the transplantation of ADMSCs is an effective method to prevent ALF in CCl_4 -induced animal model. Furthermore,

the homing of ADMSCs to the injured liver and their differentiation into hepatocytes were investigated.

2. Materials and Methods

2.1. Isolation and Culture of ADMSCs. ADMSCs were isolated from inguinal fat pads of enhanced green fluorescent protein (EGFP) transgenic C57BL/6 mice (Cyagen Biosciences, Guangzhou, China). Adipose tissues were washed extensively with phosphate-buffered saline (PBS) and digested with 0.075% collagenase type I prepared in PBS at 37°C for 60 min with shaking. Enzyme activity was neutralized by adding 5 mL of alpha-MEM containing 10% heat inactivated fetal bovine serum (FBS, Gibco, USA). The solution was filtered through a 100 μ m nylon mesh and centrifuged at 1500 rpm for 8 min. The pellet was washed twice with PBS and centrifuged at 1000 rpm for 5 min. The cells were resuspended in stromal medium (alpha-MEM, Gibco, USA) supplemented with 10% FBS and maintained in a humidified tissue culture incubator at 37°C with 5% CO₂. Upon reaching an 80% confluence, the cells were digested with 0.25% trypsin/EDTA, centrifuged, and resuspended in the same culture medium. After 3 passages, the cells were used for analysis.

2.2. Phenotypic Analysis via Flow Cytometry. Phenotypic analyses of cultured ADMSCs were performed prior to transplantation via standard flow cytometry methods. The third and fifth passage cells were harvested, suspended at a concentration of 1×10^6 cells/100 μ L in PBS, and incubated with anti-mouse monoclonal antibodies against CD44-phycoerythrin (PE), CD105-PE, CD34-allophycocyanin (APC), and CD45-APC (BD Biosciences, San Jose, CA) for 30 min in the dark at 4°C. Cells were washed and analyzed using a FACSCalibur (BD, USA).

2.3. Adipogenic and Osteogenic Differentiation. ADMSCs were plated at a density of 2×10^3 /well in six-well plates (Corning, USA) and cultured in special conditional adipogenic medium (OriCell MSC Adipogenic Differentiation Medium, Cyagen Biosciences, Guangzhou, China). After 2-3 weeks, adipogenic differentiation was evaluated by Oil red O staining. To induce osteogenic differentiation, ADMSCs were cultured in special conditional osteogenic differentiation medium (OriCell MSC Osteogenic Differentiation Medium, Cyagen Biosciences, Guangzhou, China). After 3 weeks, osteogenic differentiation was evaluated by Alizarin Red staining.

2.4. Mice Model and Cell Transplantation. Male C57BL/6 mice weighing 20–23 g were obtained from Guangdong Laboratory Animals (Guangzhou, Guangdong, China). Animal procedures were approved by the Animal Ethics Committee of Southern Medical University. To establish the acute liver failure (ALF) mice model, CCl₄ (Sigma-Aldrich) was dissolved in olive oil at 20% concentration and injected intraperitoneally into adult male C57BL/6 mice at a dose of 2.8 mL/kg. Cells were transplanted under ethyl-ether inhalation anesthesia at 24 h after administration of CCl₄.

Experimental mice were divided into two groups: transplantation group ($n = 8$), which received EGFP-ADMSCs (5×10^6 cells in 100 μ L PBS) via the tail vein; control group ($n = 6$), which receive the equal volume of PBS without cells. After the cell transplantation, blood samples were collected, and the entire livers were taken, fixed, and prepared for further analysis from sacrificed mice.

2.5. Long-Term Survival and Biochemical Analysis. Mice that had lived for more than 168 h after transplantation were considered to be survivors. Serum aspartate aminotransferase (AST) and alanine aminotransferase (ALT) levels in mice blood were analyzed prior to ADMSCs transplantation and then on days 7 and 14 after transplantation with an automated biochemical analyzer.

2.6. Pathology and Immunofluorescence. Liver tissue samples were harvested at the indicated time points after cell transplantation, fixed in 4% paraformaldehyde, embedded in paraffin, and stained with hematoxylin and eosin (H&E). Frozen sections (5 μ m thick) were used for immunofluorescence staining. Sections were blocked with 5% bovine serum albumin in PBS and then incubated with anti-mouse ALB antibody (1:200, Abcam) and anti-mouse CK18 antibody (1:250, Abcam) according to the manufacturer's instructions. Cy3-conjugated IgG antibody was used as a secondary antibody (1:500, Abcam). The sections were observed under a fluorescent microscope (OLYMPUS, BX53F, Japan).

2.7. Statistical Analysis. Survival was analyzed using a Kaplan-Meier plot and log-rank analysis. Data were presented as the mean \pm SD. Differences in serum levels of biochemical parameters were analyzed using Student's *t*-test with SPSS software version 13.0 (SPSS, Chicago, IL). The significance for all statistical analyses was defined as $P < 0.05$.

3. Results

3.1. Characteristics of ADMSCs. The morphology of transgenic ADMSCs is consistent with that of normal ADMSCs. EGFP-ADMSCs are fibroblastic in appearance and fluoresce when examined through a FITC filter (Figures 1(A) and 1(B)). The analysis by flow cytometry revealed that these cells were negative for the expression of hematopoietic markers CD45 and CD34, but positive for CD105 and CD44, which are generally considered as markers of MSCs (Figure 1(b)). The multiple differentiation potential of ADMSCs was demonstrated by inducing adipogenic and osteogenic differentiation. The adipogenic differentiation of the ADMSCs was characterized by Oil red O staining, and lipid droplets were visible in the differentiated adipocytes on day 14 after the induction of differentiation (Figure 1(C)). Osteogenic differentiation yielded an extracellular precipitate, which was identified as calcium deposits by the Alizarin Red stain; differentiated ADMSCs formed bone nodules and stained positive (Figure 1(D)). These results indicate that the cells used for transplantation exhibited the classic ADMSC phenotype and multipotential




FIGURE 1: Characteristics of ADMSCs. (a) The morphology of adipose derived mesenchymal stem cells (ADMSCs) and ADMSCs differentiated into adipocytes, osteocytes. ADMSCs exhibited a fibroblast-like morphology on light microscopic images and show green on fluorescence microscopic images (magnification $\times 10$) (A, B). The adipogenic differentiation of the ADMSCs was characterized by Oil red O (magnification $\times 20$) (C). The osteogenic differentiation of ADMSCs was shown by calcium mineralization as revealed by Alizarin Red (magnification $\times 20$) (D). (b) Phenotype profile of ADMSCs determined by flow cytometry.

stem cell characteristics, and the presence of EGFP does not seem to change stem cell properties.

3.2. Improve Survival Rate and Liver Function by ADMSCs Transplantation. The mice that survived for 7 days (168 h) could usually survive for long term. Five of six animals in a negative control group died between 12 h and 48 h after PBS treatment. However, six of eight animals treated with ADMSCs recovered from ALF. The survival rate in the ADMSCs group was significantly higher than the PBS group ($P < 0.05$) (Figure 2(a)).

As shown in Figures 2(b) and 2(c), the serum ALT and AST levels in the injured mice increased to 2800 and 2600 IU, respectively, at 24 h after administration of CCl_4 , confirming the infliction of acute extensive liver injury by CCl_4 . The mean AST and ALT levels were dramatically decreased in the ADMSCs group at 7 days after transplantation and maintained at normal levels at 14 days (Figures 2(b) and 2(c)). Compared with the control procedures, ADMSC transplantation via the tail vein significantly improved liver function and prevented death from ALF.

3.3. Promote Liver Regeneration by ADMSCs Transplantation. ALF was validated by the observation of massive necrosis during histological examination. H&E staining (Figure 3(b)) showed large areas of inflammation, sinusoid congestion and hemorrhage, and extensive hepatocyte necrosis. Very few residual hepatocytes were found and the hepatocytes present had a swollen cytoplasm. Transplantation of ADMSCs via the tail vein had a beneficial effect on recovery of the CCl_4 -injured liver. New lobules were regenerated in the surviving

animals at 7 days after infusion of ADMSC (Figure 3(c)). Microthrombosis in the sinusoids disappeared and hepatocyte necrosis was suppressed. Furthermore, the impaired hepatic lobule returned to a nearly normal architecture at 28 days after transplantation (Figure 3(d)).

3.4. Tracing of Transplanted Cells. To investigate whether EGFP-ADMSCs are capable of engrafting in CCl_4 injured liver, animals were sacrificed 1, 2, and 4 weeks after treatment. In the EGFP-ADMSCs group, the EGFP-positive cells (ADMSCs) were easily detected in the liver by fluorescence microscopy. The transplanted cells appeared in hepatic lobules as scattered individual cells at 7 days (Figure 4(b)), as small cluster at 2 weeks (Figure 4(c)), and as a mass at 4 weeks (Figure 4(d)). The cells showed a hepatocyte-like shape at 4 weeks *in vivo*. This result suggests that the transplanted cells were well integrated in the liver parenchyma. No EGFP-positive cells were observed in the liver specimens of control group (Figure 4(a)).

Four weeks after cell transplantation, the transplanted EGFP-positive cells seen in mouse liver sections were found to express cytokeratin-18 (CK-18) (Figure 5(a)) and albumin (ALB) (Figure 5(b)). Thus, the data demonstrate the capability of ADMSCs to differentiate into hepatocyte-like cells and repopulate the liver in ALF.

4. Discussion

Several reports have demonstrated the safety and promising beneficial effects of MSCs in the treatment of chronic injury [18–20] and hepatic failure [21–23]. However, the value of




FIGURE 2: Survival rate and liver function. (a) The survival rate of transplantation group (ADMSCs) was prolonged significantly compared with the control group (PBS) ($P < 0.05$). (b) Alanine aminotransferase (ALT) and (c) aspartate aminotransferase (AST) enzyme release levels in peripheral blood samples collected at pretransplantation and 7 and 14 days after transplantation. ** $P < 0.01$, versus 7 days after transplantation and before transplantation; * $P < 0.05$, versus 14 days and 7 days after transplantation and before transplantation.

ADMSCs in acute liver failure has not been studied as other sources of MSCs even though it has clinical advantages [24]. In this study, we observed that ADMSC transplantation is an effective treatment for ALF. We also showed the hepatocyte differentiation and localization of EGFP-positive cells (ADMSCs) after transplantation.

MSCs from EGFP transgenic mice are similar in their fibroblastic morphology and in differentiation capability to those from nontransgenic mice, the expression of EGFP does not seem to affect the ability to differentiate into different lineages, and cells retain their fluorescence even after undergoing differentiation [25]. Our data revealed that, besides having adipogenic and osteogenic differentiation potential *in vitro*, EGFP-ADMSCs expressed CD44 and CD105 but not hematopoietic cell markers CD45 and CD34. Furthermore, MSCs with a fluorescent marker can be used in experiments utilizing transplantation procedures since they can be easily tracked and identified. Previous study demonstrated that the fluorescently labelled MSCs bound

preferentially to injured liver when they were infused into carbon-tetrachloride injured mice [26, 27]. Additionally, MSCs represent an advantageous cell type for allogeneic transplantation since MSCs are immune privileged, therefore reducing the risks of transplant rejection [28]. We transplanted allogeneic EGFP-ADMSCs via tail vein and found that the transplanted ADMSCs then progressively migrated into the liver lobules as scattered individual cells. Although there was a decrease in EGFP-positive cells after two weeks, we found that the EGFP-positive cells proliferated and became well integrated into the liver parenchyma after four weeks. Thus, direct transplantation of ADMSCs is a suitable delivery approach for ADMSCs to reach the injury site. And direct transfusion within this damaged environment may result in the proliferation.

To date, only a few clinical studies of bone marrow stem cells transplantation for treatment of hepatic failure have been reported in the literature with limited degrees of success [29–31]. To explore the clinical value of ADMSCs,




FIGURE 3: H&E staining. (a) Normal liver tissue. (b) ALF mice showed large areas of inflammation, sinusoid congestion and hemorrhage, and extensive hepatocyte necrosis. (c) New lobules were regenerated at 7 days after infusion of ADMSCs. (d) The impaired hepatic lobule returned to a nearly normal architecture at 28 days after transplantation. Bar = 50 μm .




FIGURE 4: The fluorescent images of EGFP-ADMSCs transplanted mice liver sections. (a) Control group. The EGFP-positive cells were widely distributed in the hepatic lobules on day 7 (b), appearing as small cluster on day 14 (c) and as a mass on day 28 (d) after cell transplantation. Bar = 50 μm .




FIGURE 5: Immunofluorescence of liver sections at 28 days after cell transplantation. The EGFP-positive cell showed a hepatocyte-like shape and expressed CK-18 (a), ALB (b). Bar = 50 μ m.

we established an acute liver failure model by CCl_4 administration. Liver of ALF in animals model undergoes massive necrosis with hemorrhages involving entire lobules, which results in death within 3 days following the induction. In our study, it was found that transplantation of as few as 5×10^6 ADMSCs effectively rescued 80% of the recipient mice from ALF. The recipient mortality was apparently reduced by ADMSC treatment compared with the PBS control group. Additionally, the concentrations of ALT and AST, which are two important indicators of liver injury, were significantly decreased after ADMSCs transplantation. Histopathological evaluation of liver tissue after cell transplantation provided initial insights into the cellular targets of therapy. A striking recovery of liver structure was seen after transplantation, suggesting that ADMSCs may reconstitute liver tissue by either differentiating into hepatocytes or promoting endogenous host regeneration. These results confirmed that the direct transplantation of native ADMSCs can rescue ALF and repopulate livers of mice as an efficient alternative source.

It has been suggested that liver regeneration can occur via the transdifferentiation of the transplanted cells that reside in the regenerating tissue or cell-cell fusion [32–35]. Recently, Li et al. demonstrated that immediate intraportal transplantation of human bone marrow mesenchymal stem cells (hBMSCs) may transdifferentiate into hepatocytes during the initial stage of fulminant hepatic failure (FHF) [32]. However, other reports demonstrated that hepatogenic differentiation

after transplantation of BMSCs occurred at a low frequency in a model of acute liver injury [18, 36]. In this study, in liver sections EGFP-ADMSCs presented a liver epithelial morphology and expressed hepatocyte-specific markers as CK-18 and ALB after four weeks of administration of ADMSCs in CCl_4 -treated mice. However, we could not unequivocally prove that the transplanted ADMSCs fully differentiated into cells with functions comparable to endogenous hepatocytes and rescue the ALF animals during the initial 3 days. Kuo et al. demonstrated that the superior rescuing potential of MSCs over MDHs (mesenchymal stem cell-derived hepatocytes) as well as the ability to rescue FHF in such a short period suggested that differentiation of donor cells into hepatocytes was not the primary attribute [37]. The transplanted ADMSCs may participate in liver regeneration via proliferation and stimulate the regeneration of endogenous hepatocytes via secreted molecules rather than protected cells from necrosis.

In conclusion, our study demonstrated that ADMSCs transplantation is effective in treating CCl_4 -induced acute liver failure. ADMSCs are involved in liver regeneration *in vivo*; however, the mechanisms such as paracrine, anti-inflammatory, and antiapoptotic effects of transplanted ADMSCs remain to be resolved. In addition, although ADMSCs are an outstanding source for transplantation and offer a novel therapy for acute liver diseases, it remains to be seen whether ADMSCs transplantation will be effective for treatment of chronic liver diseases of different etiologies.

Abbreviations

ADMSCs: Adipose derived mesenchymal stem cells
 ALF: Acute liver failure
 CCl₄: Carbon tetrachloride
 EGFP: Enhanced green fluorescent protein.

Conflict of Interests

The authors have declared that no conflict of interest exists.

Acknowledgment

This work was supported by Grants from the Major Project of Chinese National Programs for Fundamental Research and Development (973 Program) (2010CB945502).

References

- [1] W. Bernal, G. Auzinger, A. Dhawan, and J. Wendon, "Acute liver failure," *The Lancet*, vol. 376, no. 9736, pp. 190–201, 2010.
- [2] B. M. Stutchfield, S. J. Forbes, and S. J. Wigmore, "Prospects for stem cell transplantation in the treatment of hepatic disease," *Liver Transplantation*, vol. 16, no. 7, pp. 827–836, 2010.
- [3] F. P. Russo and M. Parola, "Stem cells in liver failure," *Best Practice and Research: Clinical Gastroenterology*, vol. 26, no. 1, pp. 35–45, 2012.
- [4] J. D. Sipe, "Tissue engineering and reparative medicine," *Annals of the New York Academy of Sciences*, vol. 961, pp. 1–9, 2002.
- [5] K. D. Lee, T. K. C. Kuo, J. Whang-Peng et al., "In Vitro hepatic differentiation of human mesenchymal stem cells," *Hepatology*, vol. 40, no. 6, pp. 1275–1284, 2004.
- [6] M. Sgodda, H. Aurich, S. Kleist et al., "Hepatocyte differentiation of mesenchymal stem cells from rat peritoneal adipose tissue *in vitro* and *in vivo*," *Experimental Cell Research*, vol. 313, no. 13, pp. 2875–2886, 2007.
- [7] R. Anzalone, M. L. Iacono, S. Corrao et al., "New emerging potentials for human wharton's jelly mesenchymal stem cells: immunological features and hepatocyte-like differentiative capacity," *Stem Cells and Development*, vol. 19, no. 4, pp. 423–438, 2010.
- [8] Y. Fukuchi, H. Nakajima, D. Sugiyama, I. Hirose, T. Kitamura, and K. Tsuji, "Human placenta-derived cells have mesenchymal stem/progenitor cell potential," *Stem Cells*, vol. 22, no. 5, pp. 649–658, 2004.
- [9] J. M. Gimble, A. J. Katz, and B. A. Bunnell, "Adipose-derived stem cells for regenerative medicine," *Circulation Research*, vol. 100, no. 9, pp. 1249–1260, 2007.
- [10] P. A. Zuk, M. Zhu, H. Mizuno et al., "Multilineage cells from human adipose tissue: implications for cell-based therapies," *Tissue Engineering*, vol. 7, no. 2, pp. 211–228, 2001.
- [11] A. Banas, T. Teratani, Y. Yamamoto et al., "Adipose tissue-derived mesenchymal stem cells as a source of human hepatocytes," *Hepatology*, vol. 46, no. 1, pp. 219–228, 2007.
- [12] R. Yañez, M. L. Lamana, J. García-Castro, I. Colmenero, M. Ramírez, and J. A. Bueren, "Adipose tissue-derived mesenchymal stem cells have *in vivo* immunosuppressive properties applicable for the control of the graft-versus-host disease," *Stem Cells*, vol. 24, no. 11, pp. 2582–2591, 2006.
- [13] A. Banas, T. Teratani, Y. Yamamoto et al., "IFATS collection: *In vivo* therapeutic potential of human adipose tissue mesenchymal stem cells after transplantation into mice with liver injury," *Stem Cells*, vol. 26, no. 10, pp. 2705–2712, 2008.
- [14] N. Kubo, S. Narumi, H. Kijima et al., "Efficacy of adipose tissue-derived mesenchymal stem cells for fulminant hepatitis in mice induced by concanavalin A," *Journal of Gastroenterology and Hepatology*, vol. 27, no. 1, pp. 165–172, 2012.
- [15] C. K. Sun, C. L. Chang, Y. C. Lin et al., "Systemic administration of autologous adipose-derived mesenchymal stem cells alleviates hepatic ischemia-reperfusion injury in rats," *Critical Care Medicine*, vol. 40, no. 4, pp. 1279–1290, 2012.
- [16] S. J. Kim, K. C. Park, J. U. Lee, K. J. Kim, and D. G. Kim, "Therapeutic potential of adipose tissue-derived stem cells for liver failure according to the transplantation routes," *Journal of the Korean Surgical Society*, vol. 81, no. 3, pp. 176–186, 2011.
- [17] H. Aurich, M. Sgodda, P. Kaltwasser et al., "Hepatocyte differentiation of mesenchymal stem cells from human adipose tissue *in vitro* promotes hepatic integration *in vivo*," *Gut*, vol. 58, no. 4, pp. 570–581, 2009.
- [18] L. V. di Bonzo, I. Ferrero, C. Cravanzola et al., "Human mesenchymal stem cells as a two-edged sword in hepatic regenerative medicine: engraftment and hepatocyte differentiation versus profibrogenic potential," *Gut*, vol. 57, no. 2, pp. 223–231, 2008.
- [19] Y. Wang, F. Lian, J. Li et al., "Adipose derived mesenchymal stem cells transplantation via portal vein improves microcirculation and ameliorates liver fibrosis induced by CCl₄ in rats," *Journal of Translational Medicine*, vol. 10, article 133, 2012.
- [20] R. L. Pan, P. Wang, L. X. Xiang, and J. Z. Shao, "Delta-like 1 serves as a new target and contributor to liver fibrosis down-regulated by mesenchymal stem cell transplantation," *The Journal of Biological Chemistry*, vol. 286, no. 14, pp. 12340–12348, 2011.
- [21] M. Miyazaki, M. Hardjo, T. Masaka et al., "Isolation of a bone marrow-derived stem cell line with high proliferation potential and its application for preventing acute fatal liver failure," *Stem Cells*, vol. 25, no. 11, pp. 2855–2863, 2007.
- [22] S. Zhang, L. Chen, T. Liu et al., "Human umbilical cord matrix stem cells efficiently rescue acute liver failure through paracrine effects rather than hepatic differentiation," *Tissue Engineering*, vol. 18, no. 13–14, pp. 1352–1364, 2012.
- [23] H. Cao, J. Yang, J. Yu et al., "Therapeutic potential of transplanted placental mesenchymal stem cells in treating Chinese miniature pigs with acute liver failure," *BMC Medicine*, vol. 10, article 56, 2012.
- [24] T. Ochiya, Y. Yamamoto, and A. Banas, "Commitment of stem cells into functional hepatocytes," *Differentiation*, vol. 79, no. 2, pp. 65–73, 2010.
- [25] C. B. Ripoll and B. A. Bunnell, "Comparative characterization of mesenchymal stem cells from eGFP transgenic and non-transgenic mice," *BMC Cell Biology*, vol. 10, article 3, 2009.
- [26] V. Aldridge, A. Garg, N. Davies et al., "Human mesenchymal stem cells are recruited to injured liver in a β 1-integrin and CD44 dependent manner," *Hepatology*, vol. 56, no. 3, pp. 1063–1073, 2012.
- [27] D. Grisafi, M. Piccoli, M. Pozzobon et al., "High transduction efficiency of human amniotic fluid stem cells mediated by adenovirus vectors," *Stem Cells and Development*, vol. 17, no. 5, pp. 953–962, 2008.

- [28] A. Uccelli, V. Pistoia, and L. Moretta, "Mesenchymal stem cells: a new strategy for immunosuppression?" *Trends in Immunology*, vol. 28, no. 5, pp. 219–226, 2007.
- [29] L. Peng, D. Y. Xie, B. L. Lin et al., "Autologous bone marrow mesenchymal stem cell transplantation in liver failure patients caused by hepatitis B: short-term and long-term outcomes," *Hepatology*, vol. 54, no. 3, pp. 820–828, 2011.
- [30] N. Levicar, M. Pai, N. A. Habib et al., "Long-term clinical results of autologous infusion of mobilized adult bone marrow derived CD34⁺ cells in patients with chronic liver disease," *Cell Proliferation*, vol. 41, no. 1, pp. 115–125, 2008.
- [31] E. Dalakas, P. N. Newsome, S. Boyle et al., "Bone marrow stem cells contribute to alcohol liver fibrosis in humans," *Stem Cells and Development*, vol. 19, no. 9, pp. 1417–1425, 2010.
- [32] J. Li, L. Zhang, J. Xin et al., "Immediate intraportal transplantation of human bone marrow mesenchymal stem cells prevents death from fulminant hepatic failure in pigs," *Hepatology*, vol. 56, no. 3, pp. 1044–1052, 2012.
- [33] Y. Sato, H. Araki, J. Kato et al., "Human mesenchymal stem cells xenografted directly to rat liver are differentiated into human hepatocytes without fusion," *Blood*, vol. 106, no. 2, pp. 756–763, 2005.
- [34] M. H. Dahlke, R. Loi, A. Warren et al., "Immune-mediated hepatitis drives low-level fusion between hepatocytes and adult bone marrow cells," *Journal of Hepatology*, vol. 44, no. 2, pp. 334–341, 2006.
- [35] M. Miyazaki, M. Hardjo, T. Masaka et al., "Isolation of a bone marrow-derived stem cell line with high proliferation potential and its application for preventing acute fatal liver failure," *Stem Cells*, vol. 25, no. 11, pp. 2855–2863, 2007.
- [36] C. B. Rountree, X. Wang, S. Ge et al., "Bone marrow fails to differentiate into liver epithelium during murine development and regeneration," *Hepatology*, vol. 45, no. 5, pp. 1250–1260, 2007.
- [37] T. K. Kuo, S. P. Hung, C. Chuang et al., "Stem cell therapy for liver disease: parameters governing the success of using bone marrow mesenchymal stem cells," *Gastroenterology*, vol. 134, no. 7, pp. 2111.e3–2121.e3, 2008.

